

## Oscillation – Set 3

1

An oscillating block-spring system has a mechanical energy of 1.00 J, an amplitude of 10.0 cm, and a maximum speed of 1.20 m/s.

- a) What is the spring constant?
- b) What is the mass of the block?
- c) What is the frequency of oscillation?


## Oscillation – Set 3

2

Use **ENERGY** techniques to answer the following question.


A solid cylinder of mass  $M=2$  kg and radius  $R=1.0$  m is attached to a horizontal spring with spring constant  $k=100$  N/m. The cylinder can roll without slipping along the horizontal plane. When the system is displaced from the equilibrium position, it executes simple harmonic motion.

- Derive an expression for the period of the oscillations in terms of  $M$  and  $k$ .
- If the cylinder has a translational velocity of  $v_0=5.0$  m/s as it passes through equilibrium, find the phase constant, the amplitude, and the maximum acceleration of the system.



A block of mass  $m_2 = 10$  kg attached to a spring with spring constant  $k = 5$  N/m is oscillating with an amplitude of  $A = 1.5$  m horizontally on a frictionless surface. When the spring has reached its maximum extension to the left, it collides with a bullet with a mass  $m_1 = 5$  g moving  $V_b = 200$  m/s towards the right.


- What is the period of the oscillator after the collision?
- What is the amplitude of the oscillator after the collision?



## Oscillation – Set 3

4

A long uniform rod of length  $L$  and mass  $m$  is free to rotate in a *horizontal* plane about a vertical axis through its center (the picture shows a *top* view). A spring with force constant  $k$  is connected horizontally between one end of the rod and a fixed wall. When the rod is in equilibrium, it is parallel to the wall.




What is the period of the *small* oscillations that result when the rod is rotated slightly and then released?

$$I_{cm} = \frac{1}{12}ML^2 \text{ for the rod.}$$

- Use **Newton's Second Law** to find the oscillator frequency.
- Use **Energy Techniques** to find the oscillator frequency.

Use **ENERGY** techniques to answer the following question.

A block of mass  $M$  resting on a frictionless surface is attached to a stiff rod of negligible mass. The other end of the rod is attached to the top of a thin bar of length  $l$  mass  $M$  that is allowed to rotate about its center. The bottom of a bar is attached to a light spring of spring constant  $k$ . The spring is relaxed when the bar is vertical. Find the frequency of small oscillations.



Two particles are in simple harmonic motion in a straight line. They have the same amplitude and a period of 1.5 s but differ in phase by  $\pi/6$  radians.

- a) How far apart are they from one another (in terms of  $A$ ) when the lagging particle is at its maximum position?
- b) Are they moving in the same direction or opposite directions?
- c) How far apart are they 0.5 seconds later?
- d) Are they moving in the same or opposite directions then?